

CHIROPRACTIC CARE: A FLAWED RISK–BENEFIT ANALYSIS?

We have several concerns regarding Ernst's article "Chiropractic Care: Attempting a Risk–Benefit Analysis."¹

The omission of methods is most troubling. Explicit criteria should be described to prevent bias in the selection of source material. With a search of one database covering 1995–2001, we retrieved 4 additional articles on chiropractic benefits and risks (references available from the author upon request). Unspecified methods preclude other investigators from confirming or refuting the author's findings through replication, a hallmark of the scientific process.²

Regarding benefit, Ernst cited his own study³ that "revealed no compelling evidence to suggest that chiropractic yields clinical effects that are distinct from those of placebo manipulation." However, this study reviewed trials of manipulation (not "chiropractic care"), and he failed to cite another review⁴ that found manipulation to be superior to placebo for chronic pain.

Regarding risk, Ernst again cited his own study⁵ as support for his assertion that "underreporting [of complications] can be as high as 100%," but he failed to cite another article⁶ in which the author argues that overreporting of complications allegedly attributable to spinal manipulation may occur as well.

Selective reporting of results is also apparent. After 1 year follow-up in the Cherkin study, chiropractic and physical therapy patients were less disabled and were more likely to perceive their care as being very good or excellent, compared with patients receiving a treatment booklet.⁷ In the Giles trial, manipulation resulted in greater improvement in pain and disability than did acupuncture and medication.⁸ Although the clinical significance of these differences is arguable, Ernst's statement that the results do "not show an advantage of chiropractic over control treatments" is also arguable.

Ernst asserts that chiropractic "patients with low back pain often receive upper spinal manipulation." Although this may be true, failure to acknowledge this critical assertion, coupled with the use of misleading phrases such as "essentially everyone receiving chiropractic treatment is at risk," leaves little doubt as to the author's bias.

The article includes several errors. Approximately 50 000 chiropractors are in active practice in the United States,⁹ not 5000 in North America. The trials cited include patients with neck pain and sciatica, not just back pain. The visit frequency associated with an increased risk of vertebrobasilar accidents in the Rothwell study¹⁰ is "more than 2," not "more than 3."

Deficient methods, a biased sample of reference material, selective reporting of results, and prejudicial language lead us to conclude that Dr. Ernst's article is without scientific merit. The fact that his paper went through peer and editorial review and into publication is a more serious matter concerning the Journal's scientific review policies. ■

Eric L. Hurwitz, DC, PhD
William C. Meeker, DC, MPH
Monica Smith, DC, PhD

About the Authors

Eric L. Hurwitz is with the Department of Epidemiology, School of Public Health, University of California, Los Angeles, and the Southern California University of Health Sciences, Whittier. William C. Meeker and Monica Smith are with the Palmer Center for Chiropractic Research, Davenport, Iowa.

Requests for reprints should be sent to Eric L. Hurwitz, DC, PhD, UCLA School of Public Health, Department of Epidemiology, Box 951772, Los Angeles, CA 90095-1772 (email: ehurwitz@ucla.edu).

References

1. Ernst E. Chiropractic care: attempting a risk–benefit analysis. *Am J Public Health.* 2002; 92: 1603–1604.
2. Hull DL. *Science as a Process: An Evolutionary Account of the Social and Conceptual Development of Science.* Chicago: The University of Chicago Press; 1988:436.
3. Ernst E, Harkness EF. Spinal manipulation: a systematic review of sham-controlled, double-blind, randomized clinical trials. *J Pain Symptom Management.* 2001;24:879–889.
4. van Tulder MW, Koes BW, Bouter LM. Conservative treatment of acute and chronic nonspecific low back pain: a systematic review of randomized controlled trials of the most common interventions. *Spine.* 1997;22:2128–2156.
5. Ernst E. Prospective investigations into the safety

of spinal manipulation. *J Pain Symptom Management*. 2001;21:238–242.

6. Leboeuf-Yde C, Rasmussen LR, Klougart N. The risk of over-reporting spinal manipulative therapy-induced injuries: a description of some cases that failed to burden the statistics. *J Manipulative Physiol Ther*. 1996;19:536–538.

7. Cherkin DC, Deyo RA, Battie M, Street J, Barlow W. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med*. 1998;339:1021–1029.

8. Giles LGF, Muller R. Chronic spinal pain syndromes: a clinical pilot trial comparing acupuncture, a nonsteroidal anti-inflammatory drug, and spinal manipulation. *J Manipulative Physiol Ther*. 1999; 22:376–381.

9. *Occupational Outlook Handbook, 2002–2003 Edition*. U.S. Dept of Labor, Bureau of Labor Statistics; 2002:249.

10. Rothwell DM, Bondy SJ, Williams I. Chiropractic manipulation and stroke: a population-based case-control study. *Stroke*. 2001;32:1054–1060.